

# Global Patient Survey Results 2018


Mapping the Kidney Cancer Patient Experience Worldwide

## KEY FINDINGS




Total Global Participants:

**1983**




### Demographics

#### Stage of Disease




#### Participants




### Understanding of Diagnosis

Understanding their individual diagnosis, including kidney cancer subtype, helps patients evaluate treatment options, clinical trials, and long-term surveillance plans.


Yet patients reported:



did not understand their subtype




were not told their subtype at diagnosis



still did not know their subtype at time of survey

### Participation in Clinical Trials




**70%**

of patients were not asked to participate in a clinical trial.

Of those, **89%** said it was fairly likely they would have participated if asked.

### Discussion of Psychosocial Impact



of patients are impacted by psychosocial issues. Most common:




Disease-related anxiety




Fear of recurrence



Fear of dying



General anxiety



are not talking about their psychosocial issues with healthcare professionals.

**We need to talk about kidney cancer.**

For more information, visit: [www.IKCC.org/global-patient-survey](http://www.IKCC.org/global-patient-survey)



[www.ikcc.org](http://www.ikcc.org)



[@IKCCorg](https://twitter.com/IKCCorg) [@IKCCtrials](https://twitter.com/IKCCtrials)

© IKCC = International Kidney Cancer Coalition